

BIRATIONAL MODELS OF DEL PEZZO FIBRATIONS

MIKHAIL GRINENKO

1. PRELIMINARIES.

Given an algebraic variety X , we can naturally attach some objects, e.g., its field of functions $k(X)$, the essential object in birational geometry. So, assuming classification to be one of the most important problems in algebraic geometry, we may be asked to describe all algebraic varieties with the same field of functions, i.e., that are birationally isomorphic to X . Of course, "all" is a too huge class, and usually we are restricted by projective and normal varieties (though non-proper or non-normal cases may naturally arise in some questions). Typically there are two main tasks:

- A. Given a variety V , one need to determine whether it is birational to another variety W .
- B. V and W are birational to each other, and one need to get a decomposition of a birational map between them into "elementary links", i.e., birational maps that are simple enough.

The rationality problem is an essential example of task A. Recall that a variety is said to be rational if it is birational to $\mathbb{P}^n$ (or, which is the same, its field of functions is $k(x_1, \dots, x_n)$). As to task B, examples will be given below, let us only note that varieties joined by "elementary links" should belong to a more or less restrictive category (otherwise the task becomes meaningless), and such a category have to meet the following requirement: any variety can be birationally transformed to one lying in it. Minimal models and Mori fibrations in dimension 3 give examples of such categories. Then, it often happens that the indicated category, in its turn, is also too large so as to be convenient, and we outline a subcategory of "good models" (e.g., relatively minimal surfaces). Here "good" means a class of varieties that are simple enough for describing, handling, classifying, and so on. Now let us look what we have in the first three dimensions.

1.1. Curves. Normal algebraic curves are exactly smooth ones. It is well known that a birational map between projective smooth curves is an isomorphism, so the birational and biregular classifications coincide

This work was supported by the London Mathematical Society, grants RFBR no. 00-15-96085 and 02-01-00441, INTAS no. 00-0259 and 00-0269, and NWO-RFBR no. 047-008-005.

in dimension 1. Projective spaces have no moduli, so $\mathbb{P}^1$ is a unique representative of rational curves.

1.2. Surfaces. It is very well known that any birational map between smooth surfaces can be decomposed into a chain of blow-ups of points and contractions of (-1) -curves without loss of the smoothness of intermediate varieties. Now it is clear that we can successively contract (in any order) all (-1) -curves and get the so-called (relatively) minimal model (i.e., nothing to contract). So it is very convenient to use minimal models as the class of "good" models and the indicated blow-ups and contractions as elementary links. As to the rationality problem, the famous theorem of Castelnuovo says that a smooth surface X is rational if and only if $H^1(X, \mathcal{O}_X) = H^0(X, 2K_X) = 0$. This is one of the most outstanding achievements of the classical algebraic geometry.

Summarizing results in dimensions 1 and 2, we can formulate the rationality criterion as follows: X is rational if and only if all essential differential-geometric invariants, i.e., $H^0(X, (\Omega_X^p)^{\otimes m})$, vanish.

1.3. Threefolds. As soon as we get to dimension 3, the situation becomes much harder. Now it isn't obvious at all what is "a good model". We may proceed with the way as in dimension 2, i.e., contracting everything that can be contracted. This is the viewpoint of Mori's theory. But then starting from a smooth variety, we may lose smoothness very quickly and even get a "very bad" variety with two Weil divisors intersecting by a point (this is the case of "a small contraction", i.e., birational morphism which is an isomorphism in codimension 1).

Nevertheless, Mori's theory states that there is the smallest category of varieties which is stable under divisorial contractions and flips (the last is exactly a tool which allows to "correct" small contractions). In what follows we shall not need details of this theory, the reader can find them in many monographs (e.g., [21]). We only point that X belongs to the Mori category if it is a projective normal variety with at most $\mathbb{Q}$ -factorial terminal singularities. It means that every Weil divisor is $\mathbb{Q}$ -Cartier, i.e., it becomes Cartier as soon as we take it with some multiplicity, and for every resolution of singularities $\varphi : Y \rightarrow X$ we have

$$K_Y = \varphi^*(K_X) + \sum a_i E_i,$$

where all discrepancies a_i are positive rational numbers, E_i exceptional divisors, and "=" means "equal as $\mathbb{Q}$ -divisors", i.e., multiplying by a suitable integer number, we get the linear equivalency. In particular, there exists the intersection theory on such varieties, which is very similar to the usual one, the difference is mostly that we must involve rational numbers as intersection indices.

The Mori category has some nice properties. In particular, the Kodaira dimension is a birational invariant under maps in this category. We recall that the Kodaira dimension $kod(X)$ is the largest dimension

of images under (rational) maps defined by linear systems $|mK_X|$. We are mostly interested in studying varieties of negative Kodaira dimension, i.e., when $H^0(X, mK_X) = 0$ for all $m > 0$, because their birational geometry is very non-trivial.

From now on, we will only consider varieties of negative Kodaira dimension. So, what are "minimal models" in the Mori theory for such varieties? These are so-called Mori fiber spaces. By definition, a triple $\mu : X \rightarrow S$ is a Mori fiber space if X is projective, $\mathbb{Q}$ -factorial and terminal, S is a projective normal variety with $\dim S < \dim X$, and μ is an extremal contraction of fibering type, i.e., the relative Picard number $\rho(X/S) = \operatorname{rk} \operatorname{Pic}(X) - \operatorname{rk} \operatorname{Pic}(S)$ is equal to 1 and $(-K_X)$ is μ -ample.

In dimension 3 (the highest dimension where the Mori theory has been proved) we have three possible types of Mori fiber spaces (or, briefly, Mori fibrations) $\mu : X \rightarrow S$:

- 1) $\mathbb{Q}$ -Fano, if $\dim S = 0$ (i.e., S is a point);
- 2) del Pezzo fibration, if $\dim S = 1$ (the fiber over the generic point of S is a del Pezzo surface of the corresponding degree);
- 3) conic bundle, if $\dim S = 2$ (the fiber over the generic point of S is a plane conic).

In what follows, we often denote a Mori fibration $\rho : V \rightarrow S$ as V/S or even simply V , if it is clear which structure morphisms and bases are meant.

Factorization of birational maps between Mori fibrations is given by the Sarkisov program (it is proved in dimension 3, see [6]). The essential assertion of this program is the following. Suppose we have two Mori fibrations V/S and U/T and a birational map

$$\begin{array}{ccc} V & \xrightarrow{\chi} & U \\ \downarrow & & \downarrow \\ S & & T \end{array}$$

Then there exists a finite chain of birational maps

$$\begin{array}{ccccccc} X_0 & \xrightarrow{\chi_1} & X_1 & \xrightarrow{\chi_2} & X_2 & \xrightarrow{\chi_3} & \dots & \xrightarrow{\chi_{N-1}} & X_{N-1} & \xrightarrow{\chi_N} & X_N \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\ S_0 & & S_1 & & S_2 & & & & S_{N-1} & & S_N \end{array}$$

where $X_0/S_0, X_1/S_1, \dots, X_N/S_N$ are Mori fibrations, $X_0/S_0 = V/S$, $X_N/S_N = U/T$, such that $\chi = \chi_N \circ \chi_{N-1} \circ \dots \circ \chi_2 \circ \chi_1$ and any of χ_i belongs to one of the four types of elementary links listed below (see figure 1). In all cases X_1/S_1 and X_2/S_2 are Mori fibrations, ψ is an isomorphism in codimension 1 (actually, a sequence of log-flips). Then, in the case of type I, μ denotes a morphism with connected fibers, γ is an extremal divisorial contraction. Note that $\rho(S_1/S_0) = 1$. In the case of type II, γ_1 and γ_2 are extremal divisorial contractions, μ is a birational map. Type III is inverse to type I. Finally, δ_1 and δ_2 in type

IV mean morphisms with connected fibers, T is a normal variety, and $\rho(S_0/T) = \rho(S_1/T) = 1$.




FIGURE 1.

Let us note that in dimension 2 Mori fibrations are exactly relatively minimal non-singular surfaces over points or curves, isomorphisms in codimension 1 are actually biregular, and extremal divisorial contractions are blow-downs of (-1) -curves. So all elementary links are very simple. As to dimension 3, the links description just given is at bottom all we know, so up to now the Sarkisov program mostly plays a theoretical rôle.

The rationality problem for three-dimensional varieties also becomes enormously hard. During a long time, many mathematicians believed that it should be possible to find a rationality criterion more or less simple as in lower dimensions. But nearly simultaneously, in early 70th, three outstanding works of different authors which gave examples of unirational but non-rational varieties, appeared. These were Iskovskikh and Manin ([17]), Clemens and Griffiths ([4]), and Artin and Mumford ([2]). Recall that an algebraic variety is said to be unirational if there exists a rational map from projective space which is finite at the generic point. The matter is that all essential differential-geometric invariants vanish on unirational varieties as well, so we can not even hope to find something like the rationality criterion for curves and surfaces, combining these invariants. The reader may find an excellent survey of the rationality problem in higher dimensions in [15].

Nevertheless, during the last 10 years, a considerable progress in the birational classification problem in all dimension has been achieved, mostly due to the conception of birationally rigid varieties first formulated by A.V.Pukhlikov. The original version of the notion of birational rigidity is closely related to some technical features of the maximal singularities method (comparing of the canonical adjunction thresholds and all that), and still in active use. Another version arised from the Mori fibrations theory. They are not identical but coincide in many cases. In this paper we use the second one since it is more understandable.

So, first we introduce the following useful notion:

DEFINITION 1.1. *We say that Mori fibrations V/S and U/T ($\dim S = \dim T > 0$) has the same Mori structure, if there exist birational maps $\chi : V \dashrightarrow U$ and $\psi : S \dashrightarrow T$ that make the following diagram to be commutative:*

$$\begin{array}{ccc} V & \xrightarrow{\chi} & U \\ \downarrow & & \downarrow \\ S & \xrightarrow{\psi} & T \end{array}$$

In $\mathbb{Q}$ -Fano cases ($\dim S = \dim T = 0$) we say that V and U has the same Mori structure if they are biregular to each other; in other words, for any birational map $\chi : V \dashrightarrow U$ there exists a birational self-map $\mu \in \text{Bir}(V)$ such that the composition $\chi \circ \mu$ is an isomorphism.

In the sequel we will often use words "to be birational over the base" instead of "to have the same Mori structure".

Now, since it is known that any threefold X can be birationally mapped onto a Mori fibration, one can formulate the classification task as follows: to describe all Mori fibrations that are birational to X and not birational over the base to each other (i.e., have different Mori structures). Clearly, varieties X and Y are birationally isomorphic, if and only if they have the same set of Mori structures. By the way, note that only links of type II join Mori fibrations with the same structure.

In the following cases the classification becomes especially easy:

DEFINITION 1.2. *A Mori fibration V/S is said to be birationally rigid if it has a unique Mori structure (i.e., any U/T that is birational to V , is birational over the base).*

Here are some examples of (birationally) rigid and non-rigid varieties. But first let me note that any rigid variety is not rational. Indeed, $\mathbb{P}^3$ is birational to $\mathbb{P}^1 \times \mathbb{P}^2$, so we have at least three different Mori structures: $\mathbb{Q}$ -Fano ($\mathbb{P}^3$ itself), del Pezzo fibration $\mathbb{P}^1 \times \mathbb{P}^2 \rightarrow \mathbb{P}^1$, and conic bundle $\mathbb{P}^1 \times \mathbb{P}^2 \rightarrow \mathbb{P}^2$. The simplest example of birationally rigid $\mathbb{Q}$ -Fano is a smooth quartic 3-fold in $\mathbb{P}^4$, this is nothing but the result of Iskovskikh and Manin ([17]). Note that they proved even more: any birational automorphism of a smooth quartic is actually biregular (and

in general case, there are no non-trivial automorphisms at all). As to conic bundles, there exists the following result of Sarkisov ([30], [31]): any standard conic bundle $V \rightarrow S$ over a smooth rational surface S with the discriminant curve C is rigid if the linear system $|4K_S + C|$ is not empty (i.e., V/S has enough degenerations). Say, if S is a plane, it is enough for C to have degree 12 or higher. Then, a smooth cubic 3-fold in $\mathbb{P}^4$ gives us an example of non-rigid conic bundles. Indeed, the projection from a line lying on such a cubic, realizes it as a conic bundle over a plane with a quintic as the discriminant curve.

Non-singular rigid and non-rigid del Pezzo fibrations will be described in the main part of the paper. Almost all results were obtained using the maximal singularities method ([17], [25]). Here we shall not give any proves, the reader can find them in [9]–[12].

Let me conclude this section with a little remark. Minimal models or Mori fiber spaces are not convenient models all the way. In many cases, there are more preferable classes of varieties. For examples, sometimes it is useful to consider Gorenstein terminal varieties with numerically effective anticanonical divisors ([1], [7]). In other words, we have to select each time.

2. THE RIGIDITY PROBLEM FOR DEL PEZZO FIBRATIONS.

Let $\rho : V \rightarrow S$ be a Mori fibration over, S a smooth curve, η the generic point of S , and V_η the fiber over the generic point. So V_η is a non-singular del Pezzo surface of degree $d = (-K_{V_\eta})^2$ over the field of functions of S . Moreover, this is a minimal surface since $\rho(V/S) = 1$, so

$$\text{Pic}(V_\eta) \otimes \mathbb{Q} = \mathbb{Q}[-K_{V_\eta}].$$

Suppose we have another Mori fibration U/T and a birational map

$$\begin{array}{ccc} V & \xrightarrow{\chi} & U \\ \downarrow & & \downarrow \\ S & & T \end{array}$$

We would like to know as much as possible about this situation; in particular, whether χ is birational over the base or not.

In first turn, we can assume S to be rational. Indeed, suppose S is a non-rational curve. Then U can not be a $\mathbb{Q}$ -Fano threefold. The simplest reason is that the Picard group of V or any its resolution of singularities must contain a continuous part arising from S , which is impossible for any birational model of U (including U itself). Or, we can use the fact that U is rationally connected (see [22]), which is not true for V : there are no rational curves lying across the fibers of V/S (i.e., covering S).

Thus, U/T is either a del Pezzo fibration or a conic bundle. Anyway, the fibers of U/T are covered by rational curves. Again, images of these curves on V can not cover S and hence lie in the fibers of V/S . So

there exists a rational map from T to S , which is actually a morphism since the curve S is smooth, and we have the following commutative diagram:

$$\begin{array}{ccc} V & \xrightarrow{\chi} & U \\ \downarrow & & \downarrow \\ S & \xleftarrow{\psi} & T \end{array}$$

Denote ζ the generic point of T and U_ζ the fiber over the generic point. Assuming U/T to be a del Pezzo fibration and taking into account that χ is birational, we see easily that ψ is an isomorphism, so it is possible to identify η and ζ . Thus, χ induce the birational map χ_η between two del Pezzo surfaces over η . Now suppose U/T is a conic bundle. Note that we may assume χ to be a link of type I. The composition $U \rightarrow T \rightarrow S$ represents U as a fibration over S , so we can define the fiber U_η over the generic point η . It is clear that U_η is a non-singular surface fibered on conics with $\text{Pic}(U_\eta) \simeq \mathbb{Z} \oplus \mathbb{Z}$ (see the construction of the link). Moreover, we may assume U_η to be minimal. Indeed, $\rho(U_\eta/T_\eta) = 1$, so there are no (-1) -curves in the fibers of $U_\eta \rightarrow T_\eta$. Hence any (-1) -curve on U_η has to be a section (which means that, birationally, $U \rightarrow T$ is a $\mathbb{P}^1$ -bundle over T). Then simply contract any of the (-1) -curves, and you obviously get a minimal del Pezzo surface, i.e., the previous case. So, anyway, we have a birational map

$$\chi_\eta : V_\eta \dashrightarrow U_\eta$$

between two non-singular minimal surfaces defined over the field of functions of S , and U_η is either a del Pezzo surface or a conic bundle. This situation was completely studied in V.A.Iskovskikh's paper [16], theorem 2.6. In particular, if V_η is a del Pezzo surface of small degree ($d \leq 3$), then U_η is always a del Pezzo surface that is isomorphic to V_η , and χ_η is either an isomorphism if $d = 1$, or a composition of Bertini and/or Geiser involutions if $d = 2, 3$.

Thus, in what follows we assume $S = \mathbb{P}^1$. Note that the field of functions $\mathbb{C}(S)$ of S is the so-called C_1 field, and from [5] it follows that V_η always has a point over $\mathbb{C}(S)$ (i.e., $V \rightarrow S$ has a section). By the way, V_η has a Zariski dense subset of such points ([22]). Yu.I.Manin proved ([23]) that del Pezzo surfaces of degree 5 or greater with a point over a perfect field are always rational over the field itself. This is exactly our case. So if $d \geq 5$, V_η is rational over $\mathbb{C}(S) = \mathbb{C}(\mathbb{P}^1)$, hence V is rational over $\mathbb{C}$. Thus, V is birational to $\mathbb{P}^3$; in particular, V/S is always non-rigid. Now suppose $d = 4$. Then we can blow up a section of $V \rightarrow S$ and, after some log flips in the fibers, get a structure of conic bundle (or, simply, blow up a point on V_η , as in theorem 2.6 of [16]). Thus, the case $d = 4$ is non-rigid too.

So we see that, from the viewpoint of the rigidity problem, the only cases $S = \mathbb{P}^1$, $d \leq 3$, are interesting. The cases of degree 1 and 2 will

be described in details in the rest of the paper. Here we only outline some results in the case $d = 3$, which is least studied.

THEOREM 2.1 ([27]). *Let $V/\mathbb{P}^1$ be a non-singular Mori fibration on cubic surfaces with the simplest degenerations (the last means that all singular fibers have at most one ordinary double point). If 1-cycles $N(-K_V)^2 - f$ are non-effective for all $N > 0$, where f is the class of a line in a fiber, then $V/\mathbb{P}^1$ is birationally rigid.*

The indicated condition on 1-cycles (the so-called K^2 -condition) is also sufficient for degree 1 and 2 (without assumptions on degenerations) but not necessary (see examples in [9]). The requirement on degenerations arises from the technical reasons and, most probably, can be omitted. The simplest examples of non-rigid fibrations on cubic surfaces are the following.

EXAMPLE 2.2 (*Quartics with a plane*). Let $X \in \mathbb{P}^4$ be a quartic threefold containing a plane. It is easy to see that, in general case, such a quartic contains exactly 9 ordinary double points lying on the plane. Note that X is a Fano but not a $\mathbb{Q}$ -Fano variety (i.e., not in the Mori category). Indeed, hyperplane sections through the plane cut off residual cubic surfaces that intersect by these 9 singular points, which is impossible for $\mathbb{Q}$ -factorial varieties. In fact, the plane and any of these residual cubics generates the Weil divisors group of X . Now blow up the plane as a subvariety in $\mathbb{P}^4$, and let V be the strict transform of X . Then V is nothing but a Mori fibration on cubic surfaces over $\mathbb{P}^1$, and the birational morphism $V \rightarrow X$ is a small resolution with 9 lines lying over the singular points of X . It is easy to check that we can produce a flop simultaneously at these 9 lines, and then contract the strict transform of the plane, which is $\mathbb{P}^2$ with the normal bundle isomorphic to $\mathcal{O}(-2)$. We get a $\mathbb{Q}$ -Fano variety U , which has the Fano index 1 and a singular point of index 2. This U is the complete intersection of two weighted cubic hypersurfaces in $\mathbb{P}(1, 1, 1, 1, 1, 2)$. The described transformation $V/\mathbb{P}^1 \dashrightarrow U$ is nothing but a link of type 3 (ψ is the flop, γ is the contraction of $\mathbb{P}^2$, see figure 1). Conjecturally, $V/\mathbb{P}^1$ and U represent all Mori structures of X .

EXAMPLE 2.3 (*Cubic threefolds*). Let $V \in \mathbb{P}^4$ be a smooth cubic hypersurface. It is well known that V is non-rational ([4]). Nevertheless, V is very far from rigid varieties. Indeed, the projection from any line on V gives us a conic bundle over $\mathbb{P}^2$ with the discriminant curve of degree 5. Then, V is birational to a non-singular Fano variety of index 1 and degree 14, which is a section of $G(1, 5) \subset \mathbb{P}^{14}$ by a linear subspace of codimension 5. Finally, blow up any plane cubic curve on V , and you get a structure of fibration on cubic surfaces. So, V has all possible types of Mori structures in dimension 3 (i.e., $\mathbb{Q}$ -Fano, conic bundle, del Pezzo fibration).

EXAMPLE 2.4 (*Hypersurfaces of bidegree $(m, 3)$ in $\mathbb{P}^1 \times \mathbb{P}^3$*). Let $V = V_m \subset \mathbb{P}^1 \times \mathbb{P}^3$ be a non-singular hypersurface of bidegree $(m, 3)$. From the Lefschetz hyperplane sections theorem it follows that $\text{Pic}(V) \simeq \mathbb{Z}[-K_V] \oplus \mathbb{Z}[F]$, where F is the class of a fiber, so $V/\mathbb{P}^1$ is a Mori fibration on cubic surfaces by means of the projection $\mathbb{P}^1 \times \mathbb{P}^3 \rightarrow \mathbb{P}^1$. Consider the second natural projection $\pi : \mathbb{P}^1 \times \mathbb{P}^3 \rightarrow \mathbb{P}^3$. If $m = 1$ then, clearly, $\pi|_V$ gives us a birational map $V \dashrightarrow \mathbb{P}^3$, so V is rational (and, thus, non-rigid) in this case. If $m \geq 3$, then for general V the conditions of theorem 2.1 are satisfied, so $V/\mathbb{P}^1$ is rigid. We consider the most interesting case $m = 2$. First, by the result of Bardelli [3], general V_m 's are non-rational if $m \geq 2$. Now we show that $V = V_2$ is non-rigid. Note that a general $V/\mathbb{P}^1$ has exactly 27 sections that are fibers of the projection π . On the other hand, let t be any other fiber of π , then t intersects V at 2 points (not necessary different), and we can transpose these points. Thus, outside of the 27 sections, there exists an involution τ , so we have a birational self-map $\tau \in \text{Bir}(V)$. On the other hand, we can simultaneously produce a flop χ centered at the indicated sections and get another Mori fibration $U/\mathbb{P}^1$, which is also a hypersurface of bidegree $(2, 3)$ in $\mathbb{P}^1 \times \mathbb{P}^3$. This χ is a link of type IV. Note that $\dim |3(-K_V) - F| = 1$, this pencil has no base components, and the strict transforms of its elements become the fibers of $U/\mathbb{P}^1$. The same is true for the corresponding pencil on U . It only remains to notice that τ maps the pencil $|F|$ to $|3(-K_V) - F|$, so we can consider χ as the Sarkisov resolution of τ . In [32] it was shown that general V_2 have only these two Mori structure.

We conclude this section with some remarks. First, in all non-rigid cases, it is easy to check that the linear systems $|n(-K_V) - F|$ are non-empty and free from base components. In [9] the following conjecture was formulated:

CONJECTURE 2.5. *Suppose $V/\mathbb{P}^1$ is a non-singular Mori fibration on del Pezzo surfaces of degree 1, 2, or 3. Then $V/\mathbb{P}^1$ is birationally rigid, if and only if the linear systems $|n(-K_V) - F|$ are either empty or not free from base components for all $n > 0$.*

This conjecture was completely proved for degree 1 ([9]) and, under some conditions of generality for several rigid cases, for degree 2 ([9] and [10]). Moreover, for all cases that are known to be rigid or non-rigid, the statement of the conjecture holds. Finally, it does work in one direction even for a general Mori fibration on del Pezzo surfaces of degree 1:

THEOREM 2.6 ([11], theorem 3.3). *If a Mori fibration $V/\mathbb{P}^1$ on del Pezzo surfaces of degree 1 is birationally rigid, then for all $n > 0$ the linear systems $|n(-K_V) - F|$ are either empty or not free from base components.*

3. PROJECTIVE MODELS OF DEL PEZZO FIBRATIONS

In this section we construct some simple projective models for non-singular (actually, for Gorenstein) fibrations on del Pezzo surfaces of degree 1 and 2. But first let me recall the simplest projective (and weighted projective) models of del Pezzo surfaces of degree 1 and 2.

3.1. Models of del Pezzo surfaces of degree 1 and 2. Consider first the degree 1. So, let X be a non-singular del Pezzo surface of degree $d = K_X^2 = 1$. It is easy to see that a general member of the linear system $| -K_X |$ is non-singular, all its elements are irreducible and reduced curves of genus 1, and $| -K_X |$ has a unique base point P . Choose a non-singular curve $C \in | -K_X |$. Using the exact sequence

$$0 \longrightarrow \mathcal{O}_X(-(i-1)K_X) \longrightarrow \mathcal{O}_X(-iK_X) \longrightarrow \mathcal{O}_C(-iK_X) \longrightarrow 0$$

and the Kodaira vanishing theorem, we see that

$$H^0(X, -iK_X) \longrightarrow H^0(C, (-iK_X)|_C) \longrightarrow 0$$

are exact for all $i \geq 0$. So we have a surjective map of the graded algebras

$$\mathcal{A}_X = \bigoplus_{i \geq 0} H^0(X, -iK_X) \longrightarrow \mathcal{A}_C = \bigoplus_{i \geq 0} H^0(C, (-iK_X)|_C)$$

which preserves the graduation. Clearly, $\mathcal{A}_C$ is generated by elements of degree $r \leq 3$ (note that $\deg(-3K_X)|_C = 3$, i.e., $(-3K_X)|_C$ is very ample), so $\mathcal{A}_X$ is generated by elements of degree not higher than 3, too. Taking into account that

$$h^0(X, -K_X) = 2, \quad h^0(X, -2K_X) = 4, \quad h^0(X, -3K_X) = 7,$$

we can write down the generators as follows:

$$\begin{aligned} H^0(X, -K_X) &= \mathbb{C} \langle x, y \rangle, \\ H^0(X, -2K_X) &= \mathbb{C} \langle x^2, xy, y^2, z \rangle, \\ H^0(X, -3K_X) &= \mathbb{C} \langle x^3, x^2y, xy^2, y^3, zx, zy, w \rangle, \end{aligned}$$

where we agree for x and y to be elements of weight 1, z of weight 2, and w of weight 3. Thus the homogeneous components of the graded algebra $\mathcal{A}_X$ are generated by monomials $x^i y^j z^k w^l$ of the corresponding degree. Now it is clear that X can be embedded as a surface in the weighted projective space $\mathbb{P}(1, 1, 2, 3)$:

$$X = \mathbf{Proj} \mathcal{A}_X = \mathbf{Proj} \mathbb{C}[x, y, z, w]/I_X \subset \mathbb{P}(1, 1, 2, 3),$$

where I_X is a principal ideal generated by a homogeneous element of degree 6. Indeed, $h^0(X, -6K_X) = 22$, but the dimension of the homogeneous component of $\mathbb{C}[x, y, z, w]$ of degree 6 is equal to 23, so there exists exactly one linear relation in it. This relation is nothing but an equation of X . It only remains to notice that X necessarily avoids

singular points of $\mathbb{P}(1, 1, 2, 3)$, whence we can put down the equation of X as follows (stretching the coordinates if needed):

$$X = \{w^2 + z^3 + zf_4(x, y) + f_6(x, y) = 0\} \subset \mathbb{P}(1, 1, 2, 3),$$

where f_i are homogeneous polynomials of the corresponding degree.

It was the way to embed X into a very well-known threefold as a surface. However, there is another useful model of X . Let

$$\mathbb{P}(1, 1, 2, 3) \dashrightarrow \mathbb{P}(1, 1, 2)$$

be a (weighted) projection from the point $(0 : 0 : 0 : 1)$. This point does not lie on X , and as it follows from the equation of X , the restriction of the projection on X gives us a morphism of degree 2

$$\varphi : X \rightarrow \mathbb{P}(1, 1, 2).$$

We can naturally identify $\mathbb{P}(1, 1, 2)$ with a non-degenerated quadratic cone $Q \subset \mathbb{P}^3$. The base point of $|-K_X|$ lies exactly over the vertex of the cone. The ramification divisor of φ is defined by an equation of degree 3 in $Q = \mathbb{P}(1, 1, 2)$. We can also describe this picture as follows: $|-2K_X|$ has no base points and defines a morphism, which is exactly our φ . Thus, we can obtain X as a double cover of a quadratic cone in $\mathbb{P}^3$ branched along a non-singular cubic section that does not contain the vertex of the cone.

Now let us turn back to a non-singular Mori fibration $V/\mathbb{P}^1$ on del Pezzo surfaces of degree 1. Clearly, a general fiber of V is smooth. But what if $V/\mathbb{P}^1$ has degenerations? Moreover, as we will see later, it must have degenerations by the relative Pickard number arguments. So suppose X be a singular fiber of V . Then we can see that X is a projective Gorenstein (i.e., the dualizing sheaf is invertible, as it follows from the adjunction formula) Cohen-Macaulay irreducible reduced (since V is non-singular) 2-dimensional scheme, K_X is anti-ample and $K_X^2 = 1$.

First suppose X is normal. The situation was completely studied in [14], and I can simply refer the reader to it. But in order to deal with projective models of X , we only need the fact that $|-K_X|$ contains a non-singular elliptic curve ([14], proposition 4.2), and then one can repeat arguments of the non-singular case word by word. So X can be defined by a homogeneous equation of degree 6 in $\mathbb{P}(1, 1, 2, 3)$ (and as before, X avoids singular points of the weighted projective space since they are non-Gorenstein), or as a double cover of a quadratic cone in $\mathbb{P}^3$, branched along a (now singular) cubic section not passing through the vertex of the cone. As to this cubic section, it may be reducible, but all its components must be reduced. It only remains to add that the base point of $|-K_X|$ is always non-singular, and X has either only Du Val singularities, or a minimally elliptic singular point (in local coordinates defined by $r^2 + p^3 + q^6 = 0$).

The non-normal case is much harder. Nevertheless, in [28] it was shown that the projective properties of the anticanonical linear systems on X are the same as in the non-singular case. In particular, the base point of $|-K_X|$ is non-singular, X can be embedded into $\mathbb{P}(1, 1, 2, 3)$ exactly in the same way, or twice cover a quadratic cone. The unique difference is that the cubic section has to contain a non-reduced component.

We will summarize all needed results concerning to the case $d = 1$ a bit later, now let us consider del Pezzo surfaces of degree 2. This case is fairly similar to the previous one, so we only outline the key points. Consider a non-singular del Pezzo surface X , $K_X^2 = 2$. Then $|-K_X|$ is base points free and contains a non-singular element C . As before, we have a surjective map $\mathcal{A}_X \rightarrow \mathcal{A}_C$, but now these algebras are generated by elements of degree not higher than 2. We see that $h^0(X, -K_X) = 3$ and $h^0(X, -2K_X) = 7$, so suppose

$$\begin{aligned} H^0(X, -K_X) &= \mathbb{C} \langle x, y, z \rangle, \\ H^0(X, -2K_X) &= \mathbb{C} \langle x^2, y^2, z^2, xy, xz, yz, w \rangle, \end{aligned}$$

where x, y , and z are of weight 1, w is of weight 2. Then,

$$X = \mathbf{Proj} \mathcal{A}_X = \mathbf{Proj} \mathbb{C}[x, y, z, w]/I_X \subset \mathbb{P}(1, 1, 1, 2),$$

the principal ideal I_X is generated by a homogeneous element of degree 4: indeed, $h^0(X, -4K_X) = 21$, but the dimension of the homogeneous component of $\mathbb{C}[x, y, z, w]$ of degree 4 is equal to 22. Since X avoids the singular point of $\mathbb{P}(1, 1, 1, 2)$, we get the following equation of X :

$$X = \{w^2 + f_4(x, y, z) = 0\} \subset \mathbb{P}(1, 1, 1, 2),$$

where f_4 is a homogeneous polynomial of degree 4. The projection $\mathbb{P}(1, 1, 1, 2) \dashrightarrow \mathbb{P}(1, 1, 1) = \mathbb{P}^2$ from the point $(0 : 0 : 0 : 1)$, being restricted to X , gives a double cover of $\mathbb{P}^2$ branched over a non-singular quartic curve. This morphism can be also defined by the linear system $|-K_X|$.

Projective models of singular (normal and non-normal) Gorenstein del Pezzo surfaces of degree 2 have the same construction as in the non-singular case: they are defined by a quartic equation in $\mathbb{P}(1, 1, 1, 2)$ and do not contain the point $(0 : 0 : 0 : 1)$. Or, we can construct them as double coverings of $\mathbb{P}^2$ branched over quartic plane curves, but now these quartics have singular points and/or non-reduced components ([14], [28]). Briefly speaking, the situation is the same as in the case $d = 1$. In particular, normal surfaces of this type may have either only Du Val singularities, or a unique minimally elliptic singularity locally defined by an equation $r^2 + p^4 + q^4 = 0$.

We summarize the results about projective models of del Pezzo surfaces of degree 1 and 2 as follows:

PROPOSITION 3.1. *Let X be a projective Gorenstein irreducible reduced del Pezzo surface of degree 1 or 2.*

Then, in the case of degree 1, $|-K_X|$ has a unique base point, which is non-singular on X , $|-2K_X|$ is base points free, and $|-3K_X|$ is very ample and embeds X into $\mathbb{P}^6$ as a surface of degree 9. The suitable choice of the coordinates $[x, y, z, w]$ of weights $(1, 1, 2, 3)$ in $\mathbb{P}(1, 1, 2, 3)$ allows to define X by an equation of degree 6:

$$w^2 + z^3 + zf_4(x, y) + f_6(x, y) = 0,$$

where f_i are homogeneous polynomials of degree i . $|-2K_X|$ defines a finite morphism φ of degree 2:

$$\varphi : X \xrightarrow[R_Q]{2:1} Q \subset \mathbb{P}^3,$$

where Q is a non-degenerated quadratic cone, $R_Q = R|_Q$ is the ramification divisor, $R \subset \mathbb{P}^3$ is a cubic that does not pass through the vertex of Q .

In the case of degree 2, $|-K_X|$ is base points free, $|-2K_X|$ is very ample and embeds X into $\mathbb{P}^6$ as a surface of degree 8. X can be defined as a surface by an equation of degree 4:

$$w^2 + f_4(x, y, z) = 0,$$

where f_4 is homogeneous of degree 4, $[x, y, z, w]$ are the coordinates of weights $(1, 1, 1, 2)$ in $\mathbb{P}(1, 1, 1, 2)$. Finally, $|-K_X|$ defines a finite morphism of degree 2

$$\varphi : X \xrightarrow[R]{2:1} \mathbb{P}^2,$$

where R is the ramification curve, $\deg_{\mathbb{P}^2} R = 4$.

Now it is clear that models of fibrations on del Pezzo surfaces over $\mathbb{P}^1$ can be obtained as the relative version of the constructions just introduced.

3.2. Models of fibrations on del Pezzo surfaces of degree 1. Let $\rho : V \rightarrow \mathbb{P}^1$ be a non-singular fibration on del Pezzo surfaces of degree 1. Consider a sheaf of graded $\mathcal{O}_{\mathbb{P}^1}$ -algebras

$$\mathcal{A}_V = \bigoplus_{i \geq 0} \rho_* \mathcal{O}_V(-iK_V).$$

Since $(-K_V)$ is ρ -ample (i.e., on each fiber its restriction gives anti-canonical divisor, which is ample by the assumption), then clearly

$$V = \mathbf{Proj}_{\mathbb{P}^1} \mathcal{A}_V.$$

Comparing the situation with proposition 3.1, we see that there exist an algebra $\mathcal{O}_{\mathbb{P}^1}[x, y, z, w]$ of polynomials over $\mathcal{O}_{\mathbb{P}^1}$ graded according to the weights $(1, 1, 2, 3)$, and a sheaf of principal ideals $\mathcal{I}_V$ in it such that

$$V = \mathbf{Proj}_{\mathbb{P}^1} \mathcal{A}_V = \mathbf{Proj}_{\mathbb{P}^1} \mathcal{O}_{\mathbb{P}^1}[x, y, z, w]/\mathcal{I}_V \subset \mathbb{P}_{\mathbb{P}^1}(1, 1, 2, 3),$$

where $\mathbb{P}_{\mathbb{P}^1}(1, 1, 2, 3) = \mathbf{Proj}_{\mathbb{P}^1} \mathcal{O}_{\mathbb{P}^1}[x, y, z, w]$. In other words, $V/\mathbb{P}^1$ is defined by a (weighted) homogeneous polynomial (with coefficients in

$\mathcal{O}_{\mathbb{P}^1}$) in the corresponding weighted projective space over $\mathbb{P}^1$. Note that $\mathbb{P}_{\mathbb{P}^1}(1, 1, 2, 3)$ has two distinguished sections along which it is singular, so V does not intersect these sections.

Now let us construct the second model of $V/\mathbb{P}^1$. All details and proofs can be found in [9], section 2.

In what follows, we assume that

$$\text{Pic}(V) = \mathbb{Z}[-K_V] \oplus \mathbb{Z}[F],$$

where F denotes the class of a fiber. Notice that $V/\mathbb{P}^1$ has a distinguished section s_b that intersects each fiber at the base point of the anticanonical linear system. We can also define it as

$$s_b = \text{Bas} \mid -K_V + lF$$

for all $l \gg 0$. Then, $\rho_*(-2K_V + mF)$ is a vector bundle of rank 4 over $\mathbb{P}^1$. We can choose m such that

$$\mathcal{E} = \rho_*(-2K_V + mF) \simeq \mathcal{O} \oplus \mathcal{O}(n_1) \oplus \mathcal{O}(n_2) \oplus \mathcal{O}(n_3)$$

for some $0 \leq n_1 \leq n_2 \leq n_3$. Denote

$$b = n_1 + n_2 + n_3.$$

Let $X = \mathbf{Proj} \mathcal{E}$ be the corresponding $\mathbb{P}^3$ -fibration over $\mathbb{P}^1$, $\pi : X \rightarrow \mathbb{P}^1$ the natural projection. Denote M the class of the tautological bundle on X (i.e., $\pi_*\mathcal{O}(M) = \mathcal{E}$), L the class of a fiber of π , t_0 the class of a section that corresponds to the surjection $\mathcal{E} \rightarrow \mathcal{O} \rightarrow 0$, and l the class of a line in a fiber of π . Note that t_0 is nothing but a minimally twisted over the base effective irreducible section of π . Also t_0 can be obtained from the conditions $M \circ t_0 = 0$, $L \circ t_0 = 1$. So, we have described all generators of the groups of 1- and 3-dimensional cycles on X .

Again, looking at proposition 3.1, we see that there exist a threefold $Q \subset X$ fibered into quadratic cones (without degenerations) with a section t_b as the line of the cone vertices, a divisor R fibered into cubic surfaces and such that its restriction $R_Q = R|_Q$ does not intersect t_b , and a finite morphism $\varphi : V \rightarrow Q$ of degree 2 branched over R_Q , such that the following diagram becomes commutative:

$$\begin{array}{ccc} V & \xrightarrow[\substack{R_Q, 2:1}]{\varphi} & Q \subset X \\ \rho \downarrow & & \downarrow \pi = \pi|_Q \\ \mathbb{P}^1 & \xlongequal{\quad} & \mathbb{P}^1 \end{array}$$

In order to make precise the construction, suppose $t_b \sim t_0 + \varepsilon l$. Obviously, ε is some non-negative integer number.

LEMMA 3.2 ([9], lemma 2.2). *The only following cases are possible:*

- 1) $\varepsilon = 0$, i.e., $t_b = t_0$, and then $2n_2 = n_1 + n_3$, n_1 and n_3 are even, $Q \sim 2M - 2n_2L$, and $R \sim 3M$;

- 2) $\varepsilon = n_1 > 0$, and then $n_3 = 2n_2$, n_1 is even, $n_2 \geq 3n_1$, $Q \sim 2M - 2n_2L$, and $R \sim 3M - 3n_1L$.

This lemma shows that the numbers n_1 , n_2 , and n_3 completely define the construction of $V/\mathbb{P}^1$ (but up to moduli, of course). Note that these numbers are not free from relations.

By the way, we can always assume $b > 0$, i.e., $n_3 > 0$. Indeed, otherwise $Q \sim 2M$ and $\varepsilon = 0$, so V is nothing but the direct product of $\mathbb{P}^1$ and a del Pezzo surface, hence V is rational and, moreover, not a Mori fibration, because $\rho(V/\mathbb{P}^1)$ coincide with the Picard number of the del Pezzo surface in this case and thus greater than 1.

It only remains to consider some formulae and relations that allow to identify $V/\mathbb{P}^1$ more or less easy. First, let us note that a surface $G \sim (M - n_2L) \circ (M - n_3L)$ must lie in Q . This is a minimally twisted over the the base ruled surface on Q . Denote $G_V = \varphi^*(G)$ its pre-image on V . Geometrically, G_V is a minimally twisted fibration on curves of genus 1 on V . Then, t_0 lies always on Q , the fibers of Q contain lines of the class l , so the classes $s_0 = \frac{1}{2}\varphi^*(t_0)$ and $f = \frac{1}{2}\varphi^*(l)$ are well defined. Note that f is the class of the anticanonical curves in the fibers of V , and s_0 always has an effective representative in 1-cycles on V . By the way, it is easy to see that the Mori cone is generated by s_0 and f :

$$\overline{\mathbf{NE}}(V) = \mathbb{R}_+[s_0] \oplus \mathbb{R}_+[f].$$

It is not very difficult to compute the normal bundle of s_b :

$$\begin{aligned} \mathcal{N}_{s_b|V} &\simeq \mathcal{O}(-\tfrac{1}{2}n_1) \oplus \mathcal{O}(-\tfrac{1}{2}n_3), \text{ if } \varepsilon = 0, \\ \mathcal{N}_{s_b|V} &\simeq \mathcal{O}(n_1 - \tfrac{1}{2}n_3) \oplus \mathcal{O}(n_1), \text{ if } \varepsilon = n_1 > 0. \end{aligned}$$

So, in fact, we see that $V/\mathbb{P}^1$ is completely defined (again, up to moduli) by $\mathcal{N}_{s_b|V}$.

The following table contains the essential information about divisors and intersection indices on V :

$\varepsilon = 0$	$\varepsilon = n_1 > 0$
$n_1 + n_3 = 2n_2$	$n_3 = 2n_2, n_2 \geq 3n_1$
$s_b \sim s_0$	$s_b \sim s_0 + \frac{1}{2}f$
$\mathcal{N}_{s_b V} \simeq \mathcal{O}(-\frac{1}{2}n_1) \oplus \mathcal{O}(-\frac{1}{2}n_3)$	$\mathcal{N}_{s_b V} \simeq \mathcal{O}(n_1 - \frac{1}{2}n_3) \oplus \mathcal{O}(n_1)$
$K_V = -G_V + (\frac{1}{2}n_1 - 2)F$	$K_V = -G_V - (\frac{1}{2}n_1 + 2)F$
$K_V^2 = s_0 + (4 - n_2)f$	$K_V^2 = s_0 + (4 + \frac{3}{2}n_1 - n_2)f$
$s_0 \circ G_V = -\frac{1}{2}n_3$	$s_0 \circ G_V = -\frac{1}{2}n_3$
$(-K_V)^3 = 6 - 2n_2$	$(-K_V)^3 = 6 + 2n_1 - 2n_2$

REMARK 3.3. The arguments still work if we assume V to be only Gorenstein, not necessary non-singular. The only point is that we must require all fibers to be reduced. Indeed, all we need is the statement of

proposition 3.1. It is easy to see that in the Gorenstein case all fibers of $V/\mathbb{P}^1$ are irreducible (and reduced by the requirement), so we are still under the conditions even if V is singular. In particular, we have the same projective models as above, the same formulae, and so on.

3.3. Models of fibrations on del Pezzo surfaces of degree 2. Let $\rho : V \rightarrow \mathbb{P}^1$ be a non-singular fibration on del Pezzo surfaces of degree 2. As in the case $d = 1$, consider the direct image of the anticanonical algebra

$$\mathcal{A}_V = \bigoplus_{i \geq 0} \rho_* \mathcal{O}_V(-iK_V).$$

This is a sheaf of graded algebras over $\mathcal{O}_{\mathbb{P}^1}$, and $V = \mathbf{Proj}_{\mathbb{P}^1} \mathcal{A}_V$. As before, we see that there exist a polynomial algebra $\mathcal{O}_{\mathbb{P}^1}[x, y, z, w]$ graded with the weights $(1, 1, 1, 2)$, and a sheaf of principal ideals $\mathcal{I}_V$ in it such that

$$V = \mathbf{Proj}_{\mathbb{P}^1} \mathcal{A}_V = \mathbf{Proj}_{\mathbb{P}^1} \mathcal{O}_{\mathbb{P}^1}[x, y, z, w]/\mathcal{I}_V \subset \mathbb{P}_{\mathbb{P}^1}(1, 1, 1, 2).$$

Clearly, $\mathcal{I}_V$ is generated by a (weighted) homogeneous element of degree 4, and $V \subset \mathbb{P}_{\mathbb{P}^1}(1, 1, 1, 2)$ avoids a section along which $\mathbb{P}_{\mathbb{P}^1}(1, 1, 1, 2)$ is singular.

The second model is obtained by taking a double cover. We suppose $\text{Pic}(V) = \mathbb{Z}[-K_V] \oplus \mathbb{Z}[F]$. Consider

$$\mathcal{E} = \rho_* \mathcal{O}_V(-K_V + mF),$$

this is a vector bundle of rank 3 over $\mathbb{P}^1$. We may choose m such that

$$\mathcal{E} \simeq \mathcal{O} \oplus \mathcal{O}(n_1) \oplus \mathcal{O}(n_2),$$

where $0 \leq n_1 \leq n_2$. Denote $b = n_1 + n_2$.

We have the natural projection $\pi : X = \mathbf{Proj}_{\mathbb{P}^1} \mathcal{E} \rightarrow \mathbb{P}^1$. Let M denote the class of the tautological bundle, $\pi_* \mathcal{O}(M) = \mathcal{E}$, L the class of a fiber of π , l the class of a line in a fiber, and t_0 the section corresponding to the surjection $\mathcal{E} \rightarrow \mathcal{O} \rightarrow 0$, $t_0 \circ M = 0$. The following diagram is commutative:

$$\begin{array}{ccc} V & \xrightarrow[\substack{R, 2:1}]{\varphi} & X \\ \rho \downarrow & & \downarrow \pi \\ \mathbb{P}^1 & \xlongequal{\quad} & \mathbb{P}^1 \end{array}$$

Here φ is a finite morphism of degree 2 branched over a divisor $R \subset X$. R is fibered into quartic curves over $\mathbb{P}^1$, and we may suppose

$$R \sim 4M + 2aL.$$

The set (n_1, n_2, a) defines $V/\mathbb{P}^1$ up to moduli: different sets corresponds to different varieties. While n_1 and n_2 can be arbitrary chosen, a should meet some lower estimation: a can not be much less than n_1 , otherwise R does not exist.

Let us denote $H = \varphi^*(M)$, $s_0 = \frac{1}{2}\varphi^*(t_0)$, $f = \frac{1}{2}\varphi^*(l)$; clearly, $F = \varphi^*(L)$. As in the case $d = 1$, the divisor

$$G_V \sim H - n_2 F = (-K_V) + (a + n_1 - 2)F$$

plays an important role in geometry of V . This is a minimally twisted fibration into curves of genus 1 in V . Then, it is clear that

$$\overline{\mathbf{NE}}(V) = \mathbb{R}_+[s_0] \oplus \mathbb{R}_+[f],$$

but now s_0 may not correspond to an effective 1-cycle on V . The table below summarize the essential formulae on V (see [9], section 3.1):

$K_V = -H + (a + b - 2)F$
$K_V^2 = 2s_0 + (8 - 4a - 2b)f$
$(-K_V)^3 = 12 - 6a - 4b$

REMARK 3.4. We see again that all arguments and formulae work in the Gorenstein case if we keep the assumption of non-reducibility for fibers, so we may omit the non-singularity assumption. The unique difference with the case $d = 1$ is that Gorenstein singular $V/\mathbb{P}^1$ may have reducible fibers. Such fibers arise from twice covering of a plane with a double conic as the ramification divisor, so the both described models are still suitable.

4. FIBER-TO-FIBER TRANSFORMATIONS

This section is devoted to describing some special transformations of del Pezzo fibrations. The birational classification problem formulated in the bottom of section 1, is related to finding the set of all Mori structures. In other words, we study birational maps modulo birational transformations over the base. From the viewpoint of the Sarkisov program, we only need links of type I, III, and IV, i.e., only that do change the Mori structure. So links of type II, which does not change the structure of a fibration, are out of view at least for the first look. Nevertheless, it is the type of the Sarkisov links that is especially important in majority of questions related to the birational classification problem, even if we only interested in studying different Mori structures. The point is that proceeding with the maximal singularities method or, more general, the Sarkisov program and jumping from one Mori structure to another, we have to know when we should stop. So, what we really need is a criterion or at least strong enough sufficient conditions for a birational map to be over the base, i.e., fiber-to-fiber. These conditions are known (see [6], theorem 4.2(i)), but the really hard thing is to prove they are satisfied. Readers who handled any of various papers on the maximal singularities method, may remember words like "excluding infinitely near singularities", this is exactly about

the problem. Birational rigidity and birational automorphisms are important particular questions that are directly related to fiber-to-fiber transformations. Thus, it is worth to know as much as possible about this type of transformations, and this section gives some information about that.

Let V/C be a Mori fibration on del Pezzo surfaces of degree 1 or 2 over a curve C , and $\chi : V \dashrightarrow U$ a birational map onto another Mori fibration U/C that is birational over the base, i.e., we have the following commutative diagram:

$$\begin{array}{ccc} V & \xrightarrow{\chi} & U \\ \downarrow & & \downarrow \\ C & \xrightarrow{\cong} & C \end{array}$$

Taking the specialization at the generic point η of C , we see that χ induces a birational map

$$\chi_\eta : V_\eta \dashrightarrow U_\eta.$$

As it follows from [16], theorem 2.6, χ_η is an isomorphism if $d = K_{V_\eta}^2 = 1$, or decomposed into the so-called Bertini involutions if $d = 2$. Let us consider the last case in detail.

Bertini involutions can be constructed as follows. Let $A \in V_\eta$ be a rational point, i.e., defined over the field of functions of C . There exists a morphism φ of degree 2 onto a plane (proposition 3.1) branched over a (non-singular) quartic curve. Clearly φ defines an involution $\tau \in \text{Aut}(V)$ of V_η that transposes the sheets of the cover. Suppose $\varphi(A)$ does not lie on the ramification divisor, then there exists a point $A^* \in V_\eta$ which is conjugated to A by means of τ : $\tau(A) = A^*$. Clearly, $A \neq A^*$. Take a pencil of lines on the plane through $\varphi(A)$. On V_η it becomes a pencil of elliptic curves (with degenerations) with exactly two base points: A and A^* . Let us blow up these points with exceptional divisors e and e^* lying over A and A^* respectively. We obtain an elliptic surface S with 2 distinguished sections e and e^* . So, we have two (biregular) involutions μ_A and μ_{A^*} on S defined by these sections. Indeed, the specialization of S at the generic point of e (or e^* , which is the same) gives us an elliptic curve with 2 points O and O^* that corresponds to e and e^* . Each of this points can be viewed as the zero element from the viewpoint of the group law on elliptic curves, thus we get two reflection μ'_A and μ'_{A^*} defined as follows: for any point B

$$\begin{aligned} B + \mu'_A(B) &\sim 2O^*, \\ B + \mu'_{A^*}(B) &\sim 2O. \end{aligned}$$

(just in this order: μ'_A is related to O^* , μ'_{A^*} to O). These reflections give the biregular involutions μ_A and μ_{A^*} on S (note that S is relatively minimal, so any fiber-wise birational map is actually biregular). Finally, we blow down e and e^* , and then μ_A and μ_{A^*} become the desired Bertini involutions. Another way to describe μ_A is the following.

Blow up A^* , and you get a del Pezzo surface of degree 1. Its natural involution defined by the double cover of a cone (see proposition 3.1) becomes birational on V_η , and this is just our μ_A .

Thus, referring to the case $d = 2$, theorem 2.6 of [16] says: V_η is biregularly isomorphic to U_η , so we can view χ_η as a birational automorphism of V_η , and then there exists a finite set of points $I = \{A_1, A_2, \dots, A_n\}$ on V_η such that

$$\chi_\eta = \mu_{A_1} \circ \mu_{A_2} \circ \dots \circ \mu_{A_n} \circ \psi,$$

where $\psi \in \text{Aut}(V_\eta)$. Now ψ defines a fiber-wise birational transformation of $V/\mathbb{P}^1$, hence applying it to the situation, we get the case similar to degree 1, i.e., when we have an isomorphism of the fibers over the generic point.

Thus, whether $d = 1$ or $d = 2$, the picture can be reduced to the following case: we have the commutative diagram

$$(4.1) \quad \begin{array}{ccc} V & \xrightarrow{\chi} & U \\ \downarrow & & \downarrow \\ C & \xrightarrow{\simeq} & C \end{array}$$

where χ_η is an isomorphism of V_η and U_η :

$$\chi_\eta : V_\eta \xrightarrow{\simeq} U_\eta.$$

Now it is clear that if we throw out a finite number of points on C , say, $P_1, P_2, \dots, P_k$, then χ gives an isomorphism of V and U over $C \setminus \{P_1, P_2, \dots, P_k\}$. In its turn, χ can be decomposed as

$$\begin{array}{ccccccccccc} V & \xrightarrow{\chi_1} & V_1 & \xrightarrow{\chi_2} & V_2 & \xrightarrow{\chi_3} & \dots & \xrightarrow{\chi_{k-1}} & V_{k-1} & \xrightarrow{\chi_k} & U \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\ C & \xrightarrow{\simeq} & C & \xrightarrow{\simeq} & C & \xrightarrow{\simeq} & \dots & \xrightarrow{\simeq} & C & \xrightarrow{\simeq} & C \end{array}$$

where χ_i is an isomorphisms over $C \setminus \{P_i\}$. In order to distinguish such birational transformations from transformations like Bertini involution, we will call them *fiber transformations*. We may be naturally asked whether they are possible. In order to clarify the question, let us consider 2-dimensional situation. First, for any ruled surface, as an example of a fiber transformation we can take any elementary transformation of a fiber: blow up a point in a fiber and then contract the strict transform of the fiber, which is a (-1) -curve. Note that the exceptional divisor takes place of the original fiber. Obviously, such a transformation is not an isomorphism, though it gives an isomorphism of the fibers over the generic point of the base. On the other hand, consider any two relatively minimal elliptic surfaces that are birational over the base. It is very well known that such a birational map is actually an isomorphism. This is true for any relatively minimal (i.e., there are no (-1) -curves in fibers) fibrations into curves of positive genus.

What about del Pezzo fibrations? On the one hand, they are fibered into rational surfaces, and we may expect they behave like ruled surfaces. On the other hand, as it follows from results of the next two sections, their essential birational properties are far away from rational varieties, moreover, in many senses they behave like elliptic fibrations. In fact, the very situation is a bit more complicated.

The rest of this section is contained in [11], section 4 (as to the case $d = 1$), or [12] (the case $d = 2$). Let us first consider the case $d = 1$.

So, we have the commutative diagram (4.1), χ is an isomorphism of the fibers over the generic points. The situation can be easily reduced to the following case. We assume C to be a germ of a curve with the central point O , V and U Gorenstein relatively projective varieties over C fibered into del Pezzo surfaces of degree 1, their central fibers (i.e., over the point O) V_0 and U_0 reduced. Algebraically, all that means the following. Let $\mathcal{O}$ be a DVR (discrete valuation ring) with the maximal ideal $\mathfrak{m} = (t)\mathcal{O}$, where t is a local parameter, V and U non-singular del Pezzo surfaces of degree 1 defined over $\mathcal{O}$. If needed, we may take the completion of $\mathcal{O}$. Then, let K be the field of quotients, $C = \mathbf{Spec} \mathcal{O}$, $\eta = \mathbf{Spec} K$ the generic point of C . Using proposition 3.1, we may suppose that V and U are embedded into two copies P and R of the weighted projective space $\mathbb{P}_{\mathcal{O}}(1, 1, 2, 3)$ respectively. Denote $[x, y, z, w]$ and $[p, q, r, s]$ the coordinates in P and R with the weights $(1, 1, 2, 3)$.

By the condition, $\chi_{\eta} : V_{\eta} \rightarrow U_{\eta}$ is an isomorphism. The key point is that χ_{η} induces an isomorphism between P_{η} and R_{η} . It follows easily from the Kodaira vanishing theorem and the exact sequence of restriction for ideals that define V_{η} and U_{η} as surfaces in P_{η} and R_{η} respectively (see subsection 4.1, [11]).

Then, we may choose the coordinates in P and R such that

$$(4.2) \quad \begin{aligned} V &= \{w^2 + z^3 + zf_4(x, y) + f_6(x, y) = 0\} \subset P, \\ U &= \{s^2 + r^3 + rg_4(p, q) + g_6(p, q) = 0\} \subset R, \end{aligned}$$

where f_i and g_i are homogeneous polynomials of degree i . Since χ_{η} is an isomorphism of P_{η} and R_{η} , it is easy to see that χ and χ^{-1} can be defined as follows:

$$\chi = \begin{Bmatrix} p &= & t^a x \\ q &= & t^b y \\ r &= & t^c z \\ s &= & t^d w \end{Bmatrix}, \quad \chi^{-1} = \begin{Bmatrix} x &= & t^{\alpha} p \\ y &= & t^{\beta} q \\ z &= & t^{\gamma} r \\ w &= & t^{\delta} s \end{Bmatrix},$$

where each of the sets (a, b, c, d) and $(\alpha, \beta, \gamma, \delta)$ contains at least one zero. Then, all these numbers have to respect the graduation of P and R , and, moreover, we know that V and U avoid singular points of P

and R , so for some integer $m > 0$ the following conditions are satisfied:

$$\begin{cases} a + \alpha = m \\ b + \beta = m \\ c + \gamma = 2m \\ d + \delta = 3m \\ 2d = 3c \\ 2\delta = 3\gamma \end{cases}$$

Using the symmetry of the situation, we may assume that $c = 2k$, $d = 3k$, $\gamma = 2l$, $\delta = 3l$ with the conditions $k + l = m$ and $k \leq l$ (i.e., $k \leq \frac{1}{2}m$). Now substituting these relations for the numbers in (4.2), we obtain

$$(4.3) \quad \begin{aligned} f_4(x, y) &= t^{-4k} g_4(t^a x, t^b y), \\ f_6(x, y) &= t^{-6k} g_6(t^a x, t^b y). \end{aligned}$$

Suppose $k = 0$. Since the set (a, b, c, d) is not composed from zeros only (otherwise χ is already an isomorphism), then one of the numbers a and b must be positive. Let it be a . Then (4.3) shows that the equation

$$w^2 + z^3 + z f_4(x, y) + f_6(x, y) = 0$$

defines a singularity of V at the point $(t, x, y, z) = (0, 1, 0, 0, 0)$: take an affine piece $x \neq 0$ (i.e., divide the equation by x^6) and check that the differentials vanish at the indicated point.

Now let $k > 0$. Since the set (a, b, c, d) have to contain at least one zero, we may assume $a = 0$. Then $\alpha = m$. Since $l \geq k > 0$, $\alpha = m > 0$, and the set $(\alpha, \beta, \gamma, \delta)$ contains zero, we must suppose $\beta = 0$. So $b = m - \beta = m$. Thus

$$\begin{aligned} f_4(x, y) &= t^{-4k} g_4(x, t^m y), \\ f_6(x, y) &= t^{-6k} g_6(x, t^m y). \end{aligned}$$

It only remains to take into account that $k \leq \frac{1}{2}m$ and $f_i \in \mathcal{O}[x, y, z, w]$, and we see that the equation for V defines a singular point at

$$(t, x, y, z, w) = (0, 0, 1, 0, 0).$$

So we get the following assertion: V must be singular if χ is not an isomorphism!

As to the case $d = 2$ ([12]), we can repeat the previous arguments with the only difference that P and R have the type $\mathbb{P}_{\mathcal{O}}(1, 1, 1, 2)$, and V and U are defined by

$$\begin{aligned} V &= \{w^2 + f_4(x, y, z) = 0\} \subset P, \\ U &= \{s^2 + g_4(p, q, r) = 0\} \subset R. \end{aligned}$$

Then χ and χ^{-1} have the form

$$\chi = \begin{pmatrix} p &= & t^a x \\ q &= & t^b y \\ r &= & t^c z \\ s &= & t^d w \end{pmatrix}, \quad \chi^{-1} = \begin{pmatrix} x &= & t^\alpha p \\ y &= & t^\beta q \\ z &= & t^\gamma r \\ w &= & t^\delta s \end{pmatrix},$$

with the relation

$$m = a + \alpha = b + \beta = c + \gamma = \frac{1}{2}(d + \delta)$$

for some $m > 0$. Again, (a, b, c, d) and $(\alpha, \beta, \gamma, \delta)$ are not consisted of only zeros, so by the symmetry we may assume $\gamma = 0$ (thus $c = m$) and $d \geq \frac{1}{2}m$. Finally, the relation

$$f_4(x, y, z) = t^{-2d}g_4(t^ax, t^by, t^mz) \in \mathcal{O}[x, y, z]$$

shows that the equation

$$w^2 + f_4(x, y, z) = 0$$

defines a singular point at

$$(t, x, y, z, w) = (0, 0, 0, 1, 0),$$

so V is always singular if χ is not an isomorphism.

These results can be summarized as follows:

THEOREM 4.1 (Uniqueness of a smooth model). *Let V/C and U/C be non-singular Mori fibrations on del Pezzo surfaces of degree 1 or 2 that are birational over the base by means of χ :*

$$\begin{array}{ccc} V & \xrightarrow{\chi} & U \\ \downarrow & & \downarrow \\ C & \xrightarrow{\cong} & C \end{array}$$

Then V and U are always isomorphic, and χ is an isomorphism if $d = 1$, or a composition of Bertini's involutions if $d = 2$. In other words, any fiber transformation is trivial in this case.

REMARK 4.2. The similar result was proved for $d \leq 4$ in [24] using Shokurov's complements and connectedness principle. The difference is that Park's result requires V and U to have non-degenerated central fibers. In our case, we assume nothing about them.

So, the reader can see that though V/C is fibered into rational surfaces, it behaves like elliptic fibrations in dimension 2: there are no non-trivial fiber transformations without loss of smoothness. Now it is time to give some examples.

EXAMPLE 4.3 ("smooth case"). In these examples U is non-singular.

First let $d = 1$. Suppose $(a, b, c, d) = (0, 6, 2, 3)$ and $(\alpha, \beta, \gamma, \delta) = (6, 0, 10, 15)$, V and U are defined by

$$V : w^2 + z^3 + x^5y + t^{24}xy^5 = 0, \quad U : s^2 + r^3 + p^5q + pq^5 = 0.$$

It is easy to check that U is non-singular, V has a singular point of type cE_8 (the so-called compound E_8 -singularity) at $(t, x, y, z, w) = (0, 0, 1, 0, 0)$. Note that the central fiber V_0 has a unique singular point of type E_8 , U_0 is non-singular.

The case $d = 2$: suppose $(a, b, c, d) = (1, 4, 0, 2)$, $(\alpha, \beta, \gamma, \delta) = (3, 0, 4, 6)$, V and U are given by

$$V : w^2 + yz^3 + tx^4 + t^{12}y^4 = 0, \quad U : s^2 + qr^3 + tp^4 + q^4 = 0.$$

Again, U is non-singular, V has cE_8 -singularity at $(t, x, y, z, w) = (0, 0, 1, 0, 0)$. The central fiber V_0 is a non-normal del Pezzo surface (the double cover of a cone branched over a triple plane section), U_0 has an elliptic singularity.

EXAMPLE 4.4 ("birational automorphism"). The case $d = 1$: suppose $(a, b, c, d) = (1, 0, 2, 3)$, $(\alpha, \beta, \gamma, \delta) = (0, 2, 2, 3)$, and

$$V : w^2 + z^3 + t^4x^5y + xy^5 = 0, \quad U : s^2 + r^3 + p^5q + t^4pq^5 = 0.$$

V and U have cE_8 singularity in the central fibers. Note that V and U are biregularly isomorphic: put $w = s$, $z = r$, $x = q$, and $y = p$. So we can assume χ to be defined as follows:

$$x \rightarrow t^{-1}y, \quad y \rightarrow tx, \quad z \rightarrow z, \quad w \rightarrow w.$$

Thus, χ gives an example of a fiber transformation that is a birational automorphism.

The degree 2: assume $(a, b, c, d) = (1, 2, 0, 2)$, $(\alpha, \beta, \gamma, \delta) = (1, 0, 2, 2)$, V and U are given by

$$V : w^2 + t^2y^3z + yz^3 + x^4 = 0, \quad U : s^2 + q^3r + t^2qr^3 + p^4 = 0.$$

V and U have cD_4 -singularity. As before, they become isomorphic if you put $p = x$, $q = y$, $r = z$, $s = w$. So χ is a fiber transformation that is a birational automorphism.

5. MORI STRUCTURES ON DEL PEZZO FIBRATIONS: THE CASE $d = 1$

In this section we formulate known results on the rigidity of fibrations on del Pezzo surfaces of degree 1 and describe Mori structures for non-rigid cases. We use the projective model via twice covering described in subsection 3.2.

THEOREM 5.1 ([9], theorem 2.6). *Let $V/\mathbb{P}^1$ be a non-singular Mori fibration on del Pezzo surfaces of degree 1. Then $V/\mathbb{P}^1$ is birationally rigid except for two cases:*

- 1) $\varepsilon = 0$, $n_1 = n_2 = n_3 = 2$;
- 2) $\varepsilon = 0$, $n_1 = 0$, $n_2 = 1$, $n_3 = 2$.

In other words, conjecture 2.5 holds for $d = 1$. Moreover, if $V/\mathbb{P}^1$ is rigid, then this is a unique non-singular Mori fibration in its class of birational equivalency, as it follows from theorem 4.1.

REMARK 5.2. It is easy to check that linear systems $|n(-K_V) - F|$ are non-empty and free from base components exactly for the listed cases. Indeed, note that $|n(-K_V) - F|$ is non-empty and base components

free if and only if $|-K_V - F|$ has the same property (take the direct images of anticanonical linear systems), and then use the projective model. The case when $V/\mathbb{P}^1$ met the K^2 -condition was first proved in [27].

So, the only two cases are non-rigid. Let us consider them in detail.

5.1. The case $(\varepsilon, n_1, n_2, n_3) = (0, 2, 2, 2)$. First, let us note that the distinguished section s_b has the class s_0 , and this is a unique section with such a class of equivalency. Then, $\mathcal{N}_{s_b|V} \sim \mathcal{O}(-1) \oplus \mathcal{O}(-1)$, so at least locally there exists a flop centered at s_b . In fact, this flop leaves us in the projective category: it is enough to check that the linear system $n(-K_V)$ gives a birational morphism that contracts exactly s_b . Thus, let $\psi : V \dashrightarrow U$ be such a flop. Consider the linear system $\mathcal{D} = |-K_V - F|$ and its strict transform $\mathcal{D}_U = \psi_*^{-1}\mathcal{D}$ on U . It is easy to see that $\text{Bas } \mathcal{D} = s_b$ and a general member of $\mathcal{D}$ is nothing but a del Pezzo surface of degree 1 that is blown up at the base point of the anticanonical linear system. Moreover, $\dim \mathcal{D} = 1$, $\text{Bas } \mathcal{D}_U = \emptyset$, and $\mathcal{D}_U$ is a pencil of del Pezzo surfaces of degree 1. Thus, U is fibered over $\mathbb{P}^1$ onto del Pezzo surfaces of degree 1, so

$$\begin{array}{ccc} V & \xrightarrow{\psi} & U \\ \downarrow & & \downarrow \\ \mathbb{P}^1 & & \mathbb{P}^1 \end{array}$$

is a link of type IV. It only remains to compute that the projective model of $U/\mathbb{P}^1$ (as a double cover) has the same structure constants $(\varepsilon, n_1, n_2, n_3)$ as $V/\mathbb{P}^1$. We are ready to formulate the following result:

PROPOSITION 5.3 ([9], proposition 2.12). *Let $V/\mathbb{P}^1$ be a non-singular Mori fibration on del Pezzo surfaces of degree 1 with the set of structure constants $(\varepsilon, n_1, n_2, n_3) = (0, 2, 2, 2)$. Then any other Mori fibration that is birational to V , is birational over the base either to $V/\mathbb{P}^1$ itself, or to $U/\mathbb{P}^1$. $V/\mathbb{P}^1$ and $U/\mathbb{P}^1$ are the only non-singular Mori fibrations in their class of birational equivalency. In general case, $\text{Bir}(V) = \text{Aut}(V) \simeq \mathbb{Z}_2$ and generated by an automorphism corresponding to the double cover.*

REMARK 5.4. It may happen that V and U are isomorphic to each other (note they have the same set of structure constants). In this case $\psi \in \text{Bir}(V)$ and $\text{Bir}(V) \simeq \mathbb{Z}_2 \oplus \mathbb{Z}_2$.

5.2. The case $(\varepsilon, n_1, n_2, n_3) = (0, 0, 1, 2)$, or a double cone over the Veronese surface. Let $T \subset \mathbb{P}^5$ be the Veronese surface (i.e., $\mathbb{P}^2$ embedded into $\mathbb{P}^5$ by means of the complete linear system of conics), $Q \subset \mathbb{P}^6$ a cone over T with the vertex P , and $R \subset \mathbb{P}^6$ a cubic hypersurface such that $P \notin R$ and $R_Q = R \cap Q$ is non-singular. Then there exists a degree 2 finite morphism $\mu : U \rightarrow Q$ branched over R_Q . The variety U (the so-called double cone over the Veronese surface) is a

well-known Fano variety of index 2 with $\rho(U) = 1$ and $(-K_V)^3 = 8$ (see the classification of Fano 3-folds, e.g., [18]). Then, U contains a two-dimensional family $\mathcal{S}$ of elliptic curves parametrized by T . These curves lie over the rulings of Q . $\mathcal{S}$ has one-dimensional sub-family of degenerations consisting of rational curves with either a node or a cusp.

Let $l \in \mathcal{S}$ be non-singular, and $\psi_l : V_l \rightarrow U$ the blow-up of l . Then it is easy to see that V_l is a non-singular Mori fibration on del Pezzo surfaces of degree 1 with $(\varepsilon, n_1, n_2, n_3) = (0, 0, 1, 2)$. Conversely, take a del Pezzo fibration $V/\mathbb{P}^1$ with this set of structure constants. The linear system $|-K_V - 2F|$ consists of one element G_V which is the direct product of $\mathbb{P}^1$ and an elliptic curve ([9], lemma 2.9). Then the linear system $|3(-K_V) - 3F|$ defines a birational morphism $\psi : V \rightarrow U$ that contracts G_V along the rulings.

But what if $l \in \mathcal{S}$ is singular? First, suppose l is a rational curve with an ordinary double point B . Let $\psi_1 : U_1 \rightarrow U$ be the blow-up of B and $E_1 \sim \mathbb{P}^2$ the exceptional divisor. Note that l^1 (the strict transform of l) intersects E_1 at two points, and denote t_1 the line on E_1 that contains these points. Now, blow up l^1 : $\psi_2 : U_2 \rightarrow U_1$. It is easy to check that the strict transform t_1^2 of t_1 on U_2 has the normal bundle isomorphic to $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$, and we may produce a flop $\psi_3 : U_2 \dashrightarrow U_3$ centered at t_1^2 , without loss of projectivity. The strict transform E_1^3 of E_1 on U_3 becomes isomorphic to a non-singular quadric surface with the normal bundle $\mathcal{O}(-1)$, so E_1^3 can be contracted to an ordinary double point: $\psi_4 : U_3 \rightarrow U_4 = V_l$. We have the birational map $\psi_l = (\psi_4 \circ \psi_3 \circ \psi_2 \circ \psi_1)^{-1} : V_l \dashrightarrow U$, and the reader can easily check that V_l is a Gorenstein Mori fibration (over $\mathbb{P}^1$) on del Pezzo surfaces of degree 1 with a unique ordinary double point, and it has the structure set $(\varepsilon, n_1, n_2, n_3) = (0, 0, 1, 2)$ (note that the constructions in section 3 work in the Gorenstein case as well, see remark 3.3).

The case when l has a cusp at a point $B \in U$ is similar to the previous but a bit more complicated. As before, let $\psi_1 : U_1 \rightarrow U$ be the blow-up of B . We see that l^1 becomes non-singular and tangent to E_1 at some point $B_1 \in E_1$. The tangent direction to l_1 at B_1 defines a line $t_1 \subset E_1$. Now take the blow-up $\psi_2 : U_2 \rightarrow U_1$ of the curve l^1 . The strict transform E_1^2 of E_1 becomes isomorphic to a quadratic cone that is blown up at a point outside of the vertex of the cone. Moreover, the strict transform t_1^2 is exactly a unique (-1) -curve on E_1^2 and has the normal bundle $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$. All that can be checked as follows: blow up the point B_1 with an exceptional divisor E_2 , then blow up the strict transform of l_1 . The strict transform of E_2 is isomorphic to $\mathbb{F}_1$ -surface and can be contracted along its ruling. After that you get U_2 . So, now we produce a flop $\psi_3 : U_2 \dashrightarrow U_3$ centered at l_1^2 . The strict transform E_1^3 of E_1^2 is isomorphic to a quadratic cone with the normal bundle $\mathcal{O}(-1)$, hence it can be contracted $\psi_4 : U_3 \rightarrow U_4 = V_l$ to a double point locally defined by the equation $x^2 + y^2 + z^2 + w^3 = 0$.

We see that V_l is a Gorenstein Mori fibration over $\mathbb{P}^1$ on del Pezzo surfaces of degree 1 with a unique singular point, and moreover, V_l has the same structure set $(\varepsilon, n_1, n_2, n_3) = (0, 0, 1, 2)$ as before. Note that fibers of $V_l/\mathbb{P}^1$ are arised from elements of the pencil $|\frac{1}{2}(-K_V) - l|$ on U .

About twenty years ago, S.Khashin ([19]) tried to show the non-rationality of U by proving the uniqueness of the structure of a Fano variety on it. Unfortunately, his arguments contain mistakes, and up to now we have no a reliable proof of the non-rationality of U . Nevertheless, the following conjecture seems to be true:

CONJECTURE 5.5. *U has exactly the following Mori structures: U itself, and $V_l/\mathbb{P}^1$ for all $l \in \mathcal{S}$ (thus, "two-dimensional family" of different del Pezzo fibrations). In particular, U is not rational (it has no conic bundles).*

Thus, it only remains to prove this conjecture, and the birational identification problem for non-singular Mori fibrations on del Pezzo surfaces of degree 1 will be completed.

6. MORI STRUCTURES ON DEL PEZZO FIBRATIONS: THE CASE $d = 2$

We first formulate the result and then describe the non-rigid cases. The structure constants (a, n_1, n_2) are taken from subsection 3.3. Recall $b = n_1 + n_2$.

THEOREM 6.1 ([9], [10]). *Let $V/\mathbb{P}^1$ be a non-singular Mori fibration on del Pezzo surfaces of degree 2. Then $b + 2a > 0$, and if $b + 2a > 2$, $V/\mathbb{P}^1$ is birationally rigid.*

Suppose $b + 2a = 2$. Then the only following cases are possible:

- 1) $a = 0, n_1 = 0, n_2 = 2$;
- 2) $a = -2, n_1 = 2, n_2 = 4$;
- 3) $a = -3, n_1 = 2, n_2 = 6$;
- 4) $a = 1, n_1 = 0, n_2 = 0$;
- 5) $a = 0, n_1 = 1, n_2 = 1$;
- 6) $a = -1, n_1 = 2, n_2 = 2$.

General varieties in the cases 1)–3) are rigid. The cases 4)–6) are all non-rigid.

Suppose $b + 2a = 1$, then the only following cases are possible:

- 7) $a = 0, n_1 = 0, n_2 = 1$;
- 8) $a = -1, n_1 = 1, n_2 = 2$.

Both of them are non-rigid.

So, under the assumption of generality for the cases 1)–3), conjecture 2.5 holds for $d = 2$, and if $V/\mathbb{P}^1$ is rigid, then this is a unique non-singular Mori fibration in its class of birational equivalency (theorem 4.1).

REMARK 6.2. The K^2 -condition corresponds to the case $b + 2a \geq 4$, and the result was first obtained by A. Pukhlikov ([27]). The reader has not to be confused with the generality assumptions in the cases 1)–3) (see [10]), they arise from some technical troubles of the maximal singularities method. The author believes they can be omitted finally.

6.1. **The case $(a, n_1, n_2) = (1, 0, 0)$.** In this case $X \simeq \mathbb{P}^1 \times \mathbb{P}^2$, so V is nothing but a double cover of $\mathbb{P}^1 \times \mathbb{P}^2$ branched over a divisor R of bi-degree $(2, 4)$. Thus, V is a del Pezzo fibration with respect to the projection onto $\mathbb{P}^1$. On the other hand, the projection onto $\mathbb{P}^2$ represents V as a conic bundle. Indeed, a fiber of $V \rightarrow \mathbb{P}^2$ is a double cover of a line branched over 2 points, hence either a conic, or a couple of lines, or a double line. It is easy to see that the discriminant curve of $V \rightarrow \mathbb{P}^2$ has the degree 8. So, V has at least two different Mori structures. Note that $\mathbb{P}^1 \leftarrow V \rightarrow \mathbb{P}^2$ is a trivial example of type IV links.

6.2. **The case $(a, n_1, n_2) = (0, 1, 1)$, or a double quadratic cone.** In this case the linear system $|-2K_V|$ gives a small contraction onto the canonical model of V , which can be realized as double covering of a non-degenerated quadratic cone $Q \subset \mathbb{P}^4$ branched along a quartic section R_Q . It is easy to see that V has at most two curves of the class s_0 . If a curve of the class s_0 is unique on V (this means that R_Q contains the vertex of Q), then s_0 is the so-called **-2-curve** of the width 2 (in the notions of [29]). Otherwise (if R_Q does not pass through the vertex of Q), there are two curves of the class s_0 , which are disjoint and have the normal bundles $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$. In both the cases we obtain another structure of a non-singular Mori fibration on del Pezzo surfaces of degree 2 after making a flop centered at these curves. Moreover, the second structure has the same set (a, n_1, n_2) . Note that both the Mori structures arise from two families (actually, two pencils) of planes on Q .

If R_Q does not pass through the vertex of the cone, it was proved that a general variety of this type has exactly two Mori structures just described ([13]). Actually, it should be true always, not only for general cases, and even if R_Q did pass through the vertex of the cone, but the proof of this fact is still waiting for its time.

6.3. **The case $(a, n_1, n_2) = (-1, 2, 3)$.** First, let us note that V has a unique curve of the class s_0 (say, C), and it can be contracted (small contraction) by the linear system $|nH|$ for $n \geq 2$. The normal bundle of this curve is isomorphic to $\mathcal{O}(-1) \oplus \mathcal{O}(-2)$, hence there exists an anti-flip $\psi : V \dashrightarrow U$ centered at C (this is the simplest example of anti-flip, the so-called Francia's anti-flip: blow up C , make a flop centered at the minimal section of the corresponding ruled surface, and then contract the strict transform of the exceptional divisor, which is $\mathbb{P}^2$ with the normal bundle $\mathcal{O}(-2)$). U has a unique (non-Gorenstein) singular point

of index 2 (the latter means that $2K_U$ is a Cartier divisor). Moreover, U turns out to be a Mori fibration over $\mathbb{P}^1$ on del Pezzo surfaces of degree 1. It is not very difficult to check that general elements of the pencil $|-K_V - F|$ are del Pezzo surfaces of degree 1 that are blown up at the base point of the anticanonical linear system, and C is exactly their common (-1) -curve. Thus, ψ make them to be the fibers of $U/\mathbb{P}^1$. Conjecturally, $V/\mathbb{P}^1$ and $U/\mathbb{P}^1$ are the only Mori structures in this case.

6.4. The case $(a, n_1, n_2) = (0, 0, 1)$, or a double space of index 2.

This case is similar to a double cone over the Veronese surface. Let U be a double cover of $\mathbb{P}^3$ branched over a smooth quartic. This is a well known Fano variety of index 2, the so-called double space of index 2. Denote H_U the generator of the Picard group, $\text{Pic}(U) = \mathbb{Z}[H_U]$. Clearly, $K_U \sim -2H_U$. Let $l \subset U$ be a curve of genus 1 and degree 2 (i.e., $H \circ l = 2$), so l is a double cover of a line in $\mathbb{P}^3$ branched at four points. Then l is either a (non-singular) elliptic curve, or a rational curve with a double point (node or cusp), or a couple of lines with two points of intersection (probably, coincident).

Suppose l is non-singular, and $\psi_l : V_l \rightarrow U$ is the blow-up of l . Then V_l is a non-singular Mori fibration on del Pezzo surfaces of degree 2. The fibers of $V_l/\mathbb{P}^1$ arise from a pencil of planes in $\mathbb{P}^3$ that contain the image of l . It is not very difficult to compute that $V_l/\mathbb{P}^1$ has the structure set $(a, n_1, n_2) = (0, 0, 1)$. Conversely, given $V/\mathbb{P}^1$ with such a structure set, we can contract a unique divisor in $|-K_V - 2F|$, which is isomorphic to the direct product of an elliptic curve and a line, and get a double space of index 2.

If l is singular but irreducible, i.e., a rational curve with a node or a cusp, we can also obtain a Gorenstein Mori fibration on del Pezzo surfaces of degree 2, but now with a singular point of type respectively $x^2 + y^2 + z^2 + w^2 = 0$ or $x^2 + y^2 + z^2 + w^3 = 0$, "blowing up" the curve l just in the way as we obtain fibered structures from a double cone over the Veronese surface (see the previous section). $V_l/\mathbb{P}^1$ has the same set of structure constants: $(a, n_1, n_2) = (0, 0, 1)$.

However, U has another type of Mori fibration. Suppose l is a couple of lines (say, $l = l_1 \cup l_2$) intersected by two different points. Blow up one of this line and make a flop centered at the strict transform of the second line, and you obtain a structure of a (non-singular) Mori fibration on cubic surfaces $V_{l_1}/\mathbb{P}^1$. As before, its fibers arise from a pencil of planes that contain the image of any of these lines in $\mathbb{P}^3$. The same construction works if l consists of two lines that are tangent to each other. Note that we can change the order of the lines (first blowing up l_2) and get $V_{l_2}/\mathbb{P}^1$. The point is that $V_{l_1}/\mathbb{P}^1$ is (biregularly) isomorphic to $V_{l_2}/\mathbb{P}^1$ over the base, and this order change corresponds to a birational automorphism (the so-called Geiser involution, see [16]).

So we may denote these fibration on cubic surfaces $V_l/\mathbb{P}^1$ without any confusion.

Following to [20], we can formulate the following conjecture:

CONJECTURE 6.3. *Let U be a double space of index 2. Then U has the following Mori structures: U itself, and $V_l/\mathbb{P}^1$ for all curves l of degree 2 and genus 1.*

In this subsection it only remains to note that U is known to be non-rational, as it follows from [33].

6.5. The case $(a, n_1, n_2) = (-1, 1, 2)$, or a singular double cone over the Veronese surface. As to this case, note first that the linear system $|H - 2F|$ has a unique representative, which we denote G_V . Then, there is a unique curve of the class s_0 on V , and this curve has the normal bundle $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$. So we produce a flop (actually, in the projective category) $\psi_1 : V \dashrightarrow V^+$. The strict transform G_V^+ of G_V on V^+ is isomorphic to either $\mathbb{P}^1 \times \mathbb{P}^1$ or a quadratic cone, and has the normal bundle $\mathcal{O}(-1)$ in both the cases. So there exist a contraction $\psi_2 : V^+ \rightarrow U$, which gives us a Fano variety with a double point of type $x^2 + y^2 + z^2 + w^2 = 0$ or $x^2 + y^2 + z^2 + w^3 = 0$ respectively. It is easy to check that this U can be obtained as a double cover of the cone over the Veronese surface branched over a cubic section that does not pass through the vertex of the cone and has a unique du Val point of type A_1 or A_2 . Conversely, given U with such a kind of singularity, we can always obtain a structure of del Pezzo fibration as indicated. Now it is clear that U can be transformed to a (singular Gorenstein) Mori fibration on del Pezzo surfaces of degree 1 as in the previous section. Conjecturally, these are all Mori structures on U (i.e., one structure of Fano variety, one structure of fibration on del Pezzo surfaces of degree 2, and "2-dimensional family" of structures of fibrations on del Pezzo surfaces of degree 1), and hence V is a unique non-singular Mori fibration in its class of birational equivalency.

REFERENCES

- [1] Alexeev, V., *General elephants of Q -Fano 3-folds*. Compositio Math. 91 (1994), no. 1, 91–116
- [2] Artin, M., and Mumford, D., *Some elementary examples of unirational varieties which are not rational*. Proc. London Math. Soc. (3) 25 (1972), 75–95
- [3] Bardelli, F., *Polarized mixed Hodge structure: on irrationality of threefolds via degeneration*. Annali di Math. pura e appl. 137 (1984), 287–369
- [4] Clemens, H., and Griffiths, P., *The intermediate Jacobian of the cubic threefold*. Ann. of Math. (2) 95 (1972), 281–356
- [5] Colliot-Thélène, J.-L., *Arithmétique des variétés rationnelles et problèmes birationnels*. Proc. Int. Conf. Math. (1986), 641–653
- [6] Corti, A., *Factoring birational maps of threefolds after Sarkisov*. J. Algebraic Geom. 4 (1995), no. 2, 223–254
- [7] Corti, A., *Del Pezzo surfaces over Dedekind schemes*. Ann. Math., II. Ser. 144, 3 (1996), 641–683

- [8] *Flips and abundance for algebraic threefolds*. Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991. Asterisque No. 211 (1992)
- [9] Grinenko, M.M. *Birational properties of pencils of del Pezzo surfaces of degrees 1 and 2*. Sbornik: Mathematics 191, No.5, 633-653 (2000)
- [10] Grinenko, M.M. *Birational properties of pencils of del Pezzo surfaces of degrees 1 and 2. Part II*. Sbornik: Mathematics (in print)
- [11] Grinenko, M.M., *On fibrations into del Pezzo surfaces*. Math. Notes 69 (2001), no.4, 499-512
- [12] Grinenko, M.M. *On fiber-to-fiber transformations of fibrations del Pezzo surfaces of degree 2*. Russian Math. Survey 69:4 (2001)
- [13] Grinenko, M.M., *Birational automorphisms of a three-dimensional double cone*. Sbornik: Mathematics, V.189 (1998), No.7, 991-1007
- [14] Hidaka, F., Watanabe, K., *Normal Gorenstein surfaces with ample anti-canonical divisor*. Tokyo J. Math. 4 (1981), no. 2, 319-330
- [15] Iskovskikh, V.A., *On the rationality problem for algebraic threefolds*. Proc. Steklov Inst. Math. 218, 186-227 (1997)
- [16] Iskovskikh, V.A., *Factorization of birational mappings of rational surfaces from the point of view of Mori theory*. Russian Math. Surveys 51 (1996), no. 4, 585-652
- [17] Iskovskikh, V.A., and Manin, Yu.I., *Three-dimensional quartics and counterexamples to the Lüroth problem*. Math. USSR, Sbornik 15(1971), 141-166 (1972)
- [18] Iskovskikh, V.A., and Prokhorov, Yu.G., *Fano varieties*. Parshin, A.N.(ed.) et al., Algebraic Geometry V: Fano varieties. Berlin: Springer. Encycl. Math. Sci. 47, 1-245 (1999)
- [19] Khashin, S.I., *Birational automorphisms of a double Veronese cone of dimension 3*. Moscow Univ. Math. Bull., V.39 (1984), No.1, 15-20
- [20] Khashin, S.I., *Birational automorphisms of a Fano variety of index 2 and degree 2*. Moskov. Gos. Univ., manuscript deposited at VINITI, Moscow 1985. (Russian)
- [21] Kollar, J., and Mori, S., *Birational geometry of algebraic varieties*. With the collaboration of C. H. Clemens and A. Corti. Cambridge Tracts in Mathematics, 134. Cambridge University Press, Cambridge, 1998
- [22] Kollar, J., Miyaoka, Y., and Mori, S., *Rational connectedness and boundedness of Fano manifolds*. J. Differential Geom. 36 (1992), no. 3, 765-77
- [23] Manin, Ju.I., *Rational surfaces over perfect fields*. Inst. Hautes Études Sci. Publ. Math. No. 30 (1966), 55-113
- [24] Park, J., *Birational maps of del Pezzo fibrations*. J. Reine Angew. Math., V.538 (2001), 213-221
- [25] Pukhlikov, A.V., *Essentials of the method of maximal singularities*. Explicit birational geometry of 3-folds, 73-100, London Math. Soc. Lecture Note Ser., 281, Cambridge Univ. Press, Cambridge, 2000
- [26] Pukhlikov, A.V., *Birational automorphisms of Fano complete intersections*. (Russian) Dokl. Akad. Nauk 374 (2000), no. 2, 160-163
- [27] Pukhlikov, A.V., *Birational automorphisms of three-dimensional algebraic varieties with a pencil of del Pezzo surfaces*. Izv. Math. 62 (1998), no. 1, 115-155
- [28] Reid, M., *Nonnormal del Pezzo surfaces*. Publ. Res. Inst. Math. Sci. **30** (1994), No.5, 695-727
- [29] Reid, M., *Minimal models of canonical 3-folds*. Algebraic varieties and analytic varieties, Proc. Symp., Tokyo 1981, Adv. Stud. Pure Math. **1** (1983), 131-180

- [30] Sarkisov, V.G., *Birational automorphisms of conic bundles*. Math. USSR, Izv. 17, 177-202 (1981)
- [31] Sarkisov, V.G., *On conic bundle structures*. Math. USSR, Izv. 20, 355-390 (1983)
- [32] Sobolev, I.V., *Birational automorphisms of a class of varieties fibred into cubic surfaces*. Izvestiya: Mathematics, V. 66 (2002), No.1, 201-222
- [33] Tikhomirov, A.S., *The Abel-Jacobi mapping of sextics of genus 3 onto double $\mathbb{P}^3$'s of index 2*. Sov. Math., Dokl. V.33 (1986), 204-206

STEKLOV MATHEMATICAL INSTITUTE

E-mail address: grin@mi.ras.ru